

## REFERENCES

- Adelson, J.M., G.R. Helz and C.V. Miller. 2000. Reconstructing the rise of recent coastal anoxia; molybdenum in Chesapeake Bay sediments. *Geochemica et Cosmochemica Acta* 65:237-252.
- Austin, H.M., A.D. Estes and D.M. Seaver. 2000. *Estimation of juvenile striped bass: relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 1999*. Virginia Institute of Marine Science, Gloucester Point, Virginia.
- Batiuk, R.A., P. Bergstrom, M. Kemp, E. Koch, L. Murray, J.C. Stevenson, R. Bartleson, V. Carter, N.B. Rybicki, J.M. Landwehr, C. Gallegos, L. Karrh, M. Naylor, D. Wilcox, K.A. Moore, S. Ailstock and M. Teichberg. 2000. *Chesapeake Bay submerged aquatic vegetation water quality and habitat-based requirements and restoration targets: a second technical synthesis*. CBP/TRS 245/00 EPA 903-R-00-014. USEPA Chesapeake Bay Program, Annapolis, Maryland.
- Batiuk, R.A., R. Orth, K. Moore, J.C. Stevenson, W. Dennison, L. Staver, V. Carter, N. Rybicki, R. Hickman, S. Kollar and S. Bieber. 1992. *Submerged aquatic vegetation habitat requirements and restoration targets: a technical synthesis*. CBP/TRS 83/92. USEPA Chesapeake Bay Program, Annapolis, Maryland.
- Boicourt, W.C. 1992. Influences of circulation processes on dissolved oxygen in Chesapeake Bay. In Smith, D., M. Leffler and G. Mackiernan (eds.). *Oxygen dynamics in Chesapeake Bay: A synthesis of research*. University of Maryland Sea Grant College Pubs., College Park, Maryland. Pp. 7-59.
- Boreman, J. and R. Klauda. 1988. Distribution of early life stages of striped bass in the Hudson River Estuary, 1974-1979. *American Fisheries Society Monographs* 4:53-58.
- Boynton, W.R., J.H. Garber, R. Summers and W. M. Kemp. 1995. Inputs, transformations, and transport of nitrogen and phosphorous in Chesapeake Bay and selected tributaries. *Estuaries* 18:285-314.
- Bratton, J.F., S.M. Colman, R.R. Sea and P.C. Baucom. In press. Isotopic record of nitrogen and carbon cycling in Chesapeake Bay over the last 2,700 years and implications for modern oxygen depletion. *Limnology and Oceanography*.
- Bureau of Economic Analysis (BEA). 2002. Regional Economic Information System, 1969-2000. U.S. Department of Commerce, Economics and Statistics Administration.
- Bureau of Economic Analysis (BEA). 2001. Regional Economic Information System, 1969-1999. Department of Commerce, Economics and Statistics Administration.
- Carpenter, S.R. and Titus, J.E. 1984. Composition and spatial heterogeneity of submersed vegetation in a softwater lake in Wisconsin. *Vegetatio* 57:153-165.

Cerco, C.F., L. Linker, J. Sweeney, G. Shenk, and A.J. Butt. 2002. Nutrient and solids controls in Virginia's Chesapeake Bay tributaries. *Journal of Water Resources Planning and Management* May/June: 179-189.

Cerco, C.F. and K. Moore. 2001. System-wide submerged aquatic vegetation model for Chesapeake Bay. *Estuaries* 24(4): 522-534.

Cerco, C. and M. Meyers. 2000. Tributary Refinements to Chesapeake Bay Model. *Journal of Environmental Engineering* 126(2): 164-174.

Cerco, C.F. 2000. Phytoplankton kinetics in the Chesapeake Bay Eutrophication Model. *Journal of Water Quality and Ecosystem Modeling* 1(1-4): 5-49.

Cerco, C.F. 1995. Response of Chesapeake Bay to nutrient load reductions. *Journal of Environmental Engineering* 121:8 549-556.

Cerco, C.F. 1995. Simulation of Long Term Trends in Chesapeake Bay Eutrophication. *Journal of Environmental Engineering* 121(4): 298-310.

Cerco, C.F. 1993. Three-Dimensional Eutrophication Model of Chesapeake Bay. *Journal of Environmental Engineering* 119(6): 1006-1025.

Chesapeake Bay Living Resources Task Force. 1987. *Habitat requirements for Chesapeake Bay living resources: a report from the Chesapeake Bay Living Resources Task Force*. Chesapeake Bay Program, Annapolis, Maryland.

Chesapeake Bay Watershed Partners. 2001. Memorandum of understanding among the state of Delaware, the District of Columbia, the State of Maryland, the State of New York, the Commonwealth of Virginia, the State of West Virginia and the U.S. Environmental Protection Agency regarding cooperative efforts for the protection of the Chesapeake Bay and its rivers. Chesapeake Bay Program, Annapolis, Maryland.

Chesapeake Executive Council. 1987. 1987 Chesapeake Bay Agreement. Chesapeake Bay Program, Annapolis, Maryland.

Chesapeake Executive Council. 2000. Chesapeake 2000 Agreement. Chesapeake Bay Program, Annapolis, Maryland.

Colman, S.M. and J.F. Bratton. In press. Anthropogenically induced changes in sediment and biogenic silica fluxes in Chesapeake Bay. *Geology*.

Colman, S.M., P.C. Baucom, J. Bratton, T.M. Cronin, J.P. McGeehin, D.A. Willard, A. Zimmerman and P.R. Vogt. 2002. Radiocarbon dating of Holocene sediments in Chesapeake Bay. *Quaternary Research*. V. 57:58-70.

- Cooper, S.R. 1995. Chesapeake Bay watershed historical land use: Impact on water quality and diatom communities. *Ecological Applications* 5:703-723.
- Cooper, S.R. and G.S. Brush. 1991. Long-term history of Chesapeake Bay anoxia. *Science* 254:992-996.
- Cornwell, J. C., D. J. Conley, M. Owens and J. C. Stevenson. 1996. A sediment chronology of the eutrophication of Chesapeake Bay. *Estuaries* 19:488-499.
- Craig, J.K., L.B. Crowder, C.D. Gray, C.J. McDaniel, T.A. Henwood and J.G. Hanifen. 2001. Ecological effects of hypoxia on fish, sea turtles and marine mammals in the Northwestern Gulf of Mexico. Pp. 269-292. In: *Coastal Hypoxia: Consequences for Living Resources and Ecosystems*, N.N. Rabalais and R.E. Turner, eds. Coastal and Estuarine Studies.
- Cronin, T. M., ed. 2000. Initial Report on IMAGES V Cruise of the Marion-Dufresne to Chesapeake Bay June 20-22, 1999. USGS Open-file report 00-306.
- Cronin, T. M. and C. Vann. In press. The sedimentary record of anthropogenic and climatic influence on the Patuxent estuary and Chesapeake Bay ecosystems. *Estuaries*.
- Dennison, W.C., R.J. Orth, K.A. Moore, J.C. Stevenson, V. Carter, S. Kollar, P.W. Bergstrom and R.A. Batiuk. 1993. Assessing water quality with submersed aquatic vegetation. Habitat requirements as barometers of Chesapeake Bay health. *Bioscience* 43:86-94.
- Dey, W.P. 1981. Mortality and growth of young-of-the-year striped bass in the Hudson River Estuary. *Transactions of the American Fisheries Society*. 110:151-157.
- Di Toro, D. M. 2001. Sediment Flux Modeling. John Wiley and Sons, Inc. New York, 624p.
- Donigian, J., Anthony S., B. R. Bicknell, A.S. Patwardhan, L.C. Linker, C.H. Chang, R. Reynolds. 1994. Watershed Model Application to Calculate Bay Nutrient Loadings: Final Findings and Recommendations. Annapolis, MD, U.S. EPA Chesapeake Bay Program.
- Eby, L.A. 2001. Response of a fish community to frequent and infrequent disturbances in an estuarine ecosystem. Ph.D. dissertation. Duke University.
- Funderburk, S.L., J.A. Mihursky, S.J. Jordan and D. Riley (eds.). 1991. Habitat requirements for Chesapeake Bay living resources: second edition. Chesapeake Research Consortium, Solomons, Maryland.
- Hagy, J.D. 2002. Eutrophication, hypoxia and trophic transfer efficiency in Chesapeake Bay. Ph.D. dissertation, University of Maryland, College Park, Maryland.
- Harding, L.W. and E.S. Perry. 1997. Long-term increase of phytoplankton biomass in Chesapeake Bay, 1950-1994. *Marine Ecology Progress Series* 157:39-52

Howell, P. and D. Simpson. 1994. Abundance of marine resources in relation to dissolved oxygen in Long Island Sound. *Estuaries* 17:394-408.

Jordan, S.J., C. Stenger, M. Olson, R. Batiuk and K. Mountford. 1992. Chesapeake Bay dissolved oxygen goal for restoration of living resource habitats: a synthesis of living resource requirements with guidelines for their use in evaluating model results and monitoring information. CBP/TRS 88/93. Chesapeake Bay Program, Annapolis, Maryland.

Karlsen, A.W., T.M. Cronin, S. E. Ishman, D.A. Willard, R. Kerhin, C.W. Holmes and M. Marot. 2000. Historical trends in Chesapeake Bay dissolved oxygen based on benthic foraminifera from sediment cores. *Estuaries* 23:488-508.

Keister, J.E., E.D. Houde and D.L. Breitburg. 2000. Effects of bottom-layer hypoxia on abundances and depth distributions of organisms in Patuxent River, Chesapeake Bay. *Marine Ecology Progress Series* 205:43-59.

Kuenen, M.M. and A.O. Debrot. 1995. A quantitative study of the seagrass and algal meadows of the Spaanse Water, Curacao, the Netherlands, Antilles. *Aquatic Botany* 51 291-310.

Koch, E.W. 2001. Beyond light: physical, geological, and geochemical parameters as possible submersed aquatic vegetation habitat requirements. *Estuaries* 24:1-17.

Lehmann, A.E., E. Castella and J.B. Lacharanne. 1997. Morphological traits and spatial heterogeneity of aquatic plants along sediment and depth gradients, Lake Geneva, Switzerland. *Aquatic Botany* (55) 281-299.

Linker, L.C., G.W. Shenk, P. Wang, C.F. Cerco, A.J. Butt, P.J. Tango, and R.W. Savidge. 2002. A Comparison of Chesapeake Bay Estuary Model Calibration With 1985 - 1994 Observed Data and Method of Application to Water Quality Criteria, Modeling Subcommittee, Chesapeake Bay Program Office, November, 2002.

Linker, L.C., G.W. Shenk, D.L. Dennis, and J.S. Sweeney. 2000. Cross-Media Models of the Chesapeake Bay Watershed and Airshed. *Water Quality and Ecosystem Modeling* 1(1-4): 91-122.

Linker, L.C., 1996. Models of the Chesapeake Bay. *Sea Technology* 37(9):49-55.

Linker, L.C., C.G. Stigall, C.H. Chang, and A.S. Donigian, Jr., 1996. Aquatic accounting: Chesapeake Bay Watershed Model quantifies nutrient loads. *Water Environment and Technology* 8(1): 48-52.

MacGregor, J. and E.D. Houde. 1996. Onshore-offshore pattern and variability in distribution and abundance of bay anchovy *Anchoa mitchilli* eggs and larvae in Chesapeake Bay. *Marine Ecology Progress Series* 138:15-25.

Malone, T.C. 1992. Effects of water column processes on dissolved oxygen: nutrients, phytoplankton and zooplankton. In: Smith, D., M. Leffler, G. Mackiernan (eds.). Oxygen dynamics in Chesapeake Bay: A synthesis of research. University of Maryland Sea Grant College Publications., College Park, Maryland. Pp. 61-112.

Malone, T.C., W.M. Kemp, H.W. Ducklow, W.R. Boynton, J.H. Tuttle and R.B. Jonas. 1986. Lateral variation in the production and fate of phytoplankton in a partially stratified estuary. *Marine Ecology Progress Series* 32:149-160.

Manning, J. H. 1957. The Maryland Soft Shell Clam Industry and its Effects on Tidewater Resources: an Interim Report to the Maryland General Assembly. Maryland Department of Research and Education Resource Study Report No. 11. Chesapeake Biological Laboratory, Solomons, MD. 25 pages.

Mansueti, R.J. and E.H. Hollis, 1963. Striped bass in Maryland tidewater. Natural Resources Institute, University of Maryland Education Series Number 61. 28 pages.

Miller, T.J., J. Stone and E.A. Sadler. 1997. Chapter 2, Gill Net Surveys, pp. 34-63. In: Assessment of fish community structure near Site 104, T.J. Miller. (ed.) UMCES CBL 97-123. Report to the Maryland Environmental Service, Annapolis, Maryland

Moore, K., D. Wilcox, R. Orth and E. Bailey. 1999. Analysis of historical distribution of submerged aquatic vegetation (SAV) in the James River. Special Report No. 355 in *Applied Marine Science and Ocean Engineering*. Virginia Institute of Marine Science, School of Marine Science, College of William and Mary, Gloucester Point, Virginia.

Moore, K., D. Wilcox and B. Anderson. 2001. Analysis of historical distribution of submerged aquatic vegetation (SAV) in the York and Rappahannock rivers as evidence of historical water quality conditions. Special Report No. 375 in *Applied Marine Science and Ocean Engineering*. Virginia Institute of Marine Science, School of Marine Science, College of William and Mary, Gloucester Point, Virginia.

Naylor, M.D. 2002. Historic distribution of submerged aquatic vegetation (SAV) in Chesapeake Bay, Maryland. Maryland Department of Natural Resources, Annapolis, Maryland.

Newcombe, C.L. and W.A. Horne. 1938. Oxygen-poor waters of the Chesapeake Bay. *Science* 88:80-81.

Officer, C.B., R.B. Biggs, J.L. Taft, L.E. Cronin, M.A. Tyler and W.R. Boynton. 1984. Chesapeake Bay anoxia: Origin, development, and significance. *Science* 223:22-27.

Orth, R. J. and K.D. Moore. 1983. Chesapeake Bay: an unprecedented decline in submerged aquatic vegetation. *Science* 222:51-53.

Orth, R.J., D.J. Wilcox, L.S. Nagey, J.R. Whiting and J.R. Fishman. 2000. Distribution and abundance of submerged aquatic vegetation in the Chesapeake Bay and coastal bays–1999. VIMS Special Scientific Report Number 141. Final Report to the U.S. Environmental Protection Agency, Chesapeake Bay Program, Annapolis, Maryland.

Pothoven, S.A., S.B. Brandt, J.M. Jech and J.L. Horne. 1997. Chapter 3, Fish Abundance, Size and Species Composition: Kent Island Site 104, pp. 78-125. In: *Assessment of Fish Community Structure Near Site 104*. T.J. Miller (ed.). UMCES CBL 97-123. Report to the Maryland Environmental Service, Annapolis, Maryland.

Rickhus, W. and F. Stroup. 1987. Source document: characterization of the biology of and fishery of Maryland stocks of yellow perch. Maryland Department of Natural Resources Tidewater Administration, Annapolis, Maryland. Contract No. F40-85-008.

Rilling, G.C. and E.D. Houde. 1999. Regional and temporal variability in distribution and abundance of bay anchovy (*Anchoa mitchilli*) eggs and larvae in the Chesapeake Bay. *Estuaries* 22(4):1096-1109.

Secor, D.H., E. Niklitschek, J.T. Stevenson, T.E. Gunderson, S. Minkinen, B. Florence, M. Mangold, J. Skjeveland and A. Henderson-Arzapalo. 2000. Dispersal and growth of yearling Atlantic sturgeon *Acipenser oxyrinchus* released into the Chesapeake Bay. *Fisheries Bulletin* 98(4):800-810.

Settler-Hamilton, E.M., W.R. Boynton, J.A. Mihursky, T.T. Polgar and K.V. Wood. 1981. Spatial and temporal distribution of striped bass eggs, larvae and juveniles in the Potomac Estuary. *Transactions of the American Fisheries Society* 110:121-136.

Settler-Hamilton, E.M. and L. Hall, Jr. 1991. Striped bass. In: Habitat requirements for Chesapeake Bay living resources: second edition. Funderburk, S.L., J.A. Mihursky, S.J. Jordan and D. Riley (eds.). Chesapeake Research Consortium, Solomons, Maryland.

Smith, D.E., M. Leffler and G. MacKiernan. 1992. Oxygen dynamics in the Chesapeake Bay: a synthesis of recent research. Maryland and Virginia Sea Grant Programs. College Park, Maryland.

Taft, J.L., W.R. Taylor, E.O. Hartwig and R. Loftus. 1980. Seasonal oxygen depletion in Chesapeake Bay. *Estuaries* 3:242-247.

Thomann, R.V., J.R. Collier, A. Butt, E. Casman, and L.C. Linker. 1994. Response of the Chesapeake Bay Water Quality Model to Loading Scenarios. Annapolis, MD, Chesapeake Bay Program Office.

Tresselt, E.F. 1952. Spawning grounds of the striped bass, *Roccus saxatilis*, in Virginia. Bulletin of the Bingham Oceanographic Collection. 14(1): 98-110.

Tsai, C. and G. Gibson. 1971. Fecundity of the yellow perch *Perca flavescens* (Mitchil), in the Patuxent River, Maryland. *Chesapeake Science* 12:270-284.

Tuttle, J.H., R.B. Jonas and T.C. Malone. 1987. Origin, development, and significance of Chesapeake Bay anoxia. In: *Contaminant Problems and Management of Living Chesapeake Bay Resources* S.K. Majumdar, L.W. Hall, Jr. and K.M. Austin (eds.). Pennsylvania Academy of Science, Philadelphia, Pennsylvania, pp. 442-472.

U.S. Census Bureau. 2002. 1990 Census of Population and Housing. Department of Commerce, Economics, and Statistics Administration.

U.S. EPA. 2003. Ambient Water Quality Criteria for Dissolved Oxygen, Water Clarity, and Chlorophyll a for Chesapeake Bay and its Tidal Tributaries. U.S. EPA, Region 3.

U.S. Environmental Protection Agency. 2003a. Biological Evaluation for the Recommended Ambient Water Quality Criteria and Designated Uses for the Chesapeake Bay and its Tidal Waters Under the Clean Water Act Section 117. Chesapeake Bay Program Office, Annapolis, Maryland.

U.S. EPA Chesapeake Bay Program. 2002. Nutrient Reduction Technology Cost Estimations for Point Sources in the Chesapeake Bay Watershed. Chesapeake Bay Program, Annapolis, MD.

U.S. EPA. 1998. Water Quality Standards Regulation. EPA 823-Z-98-002.

U.S. EPA. 1995. Interim Economic Guidance for Water Quality Standards, Workbook. Office of Water.

U.S. EPA. 1983. Water Quality Standards Handbook. U.S. EPA, Office of Water, Office of Science and Technology, Washington D.C.

Wang, H.V. and B.H. Johnson. 2000. Validation and application of the second generation three-dimensional hydrodynamic model of Chesapeake Bay. *Journal of Water Quality and Ecosystem Modeling* 1(1-4): 51-90.

Welsh, S.A., J.E. Skjeveland, M.F. Mangold and S.M. Eylver. 2000. Distribution of wild and hatchery-reared Atlantic sturgeon in the Chesapeake Bay, Maryland. Biology, Management, and Protection of Sturgeon Symposium, Electric Power Research Institute, Palo Alto, California.

Willard, D.A., T.M. Cronin, S. Verardo. 2003. Late-Holocene climate and ecosystem history from Chesapeake Bay sediment cores, USA. *The Holocene* v. 13.

Zheng, Y., B. Weinman, T.M. Cronin, M.Q. Fleisher and R.F. Anderson. In press. A rapid procedure for thorium, uranium, cadmium and molybdenum in small sediment samples by inductively coupled plasma-mass spectrometry: Application in Chesapeake Bay. *Applied Geochemistry*.

---

Zimmerman, A.R. and E.A. Canuel. 2000. A geochemical record of eutrophication and anoxia in Chesapeake Bay sediments: Anthropogenic influence on organic matter composition. *Marine Chemistry* 69:117-137.